

# ***ina-1* expression in the somatic gonad during late larval development**


Srimoyee Ghosh<sup>1</sup>, Takao Inoue<sup>3</sup> and Paul Sternberg<sup>1,2§</sup>

<sup>1</sup>Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA

<sup>2</sup>Howard Hughes Medical Institute

<sup>3</sup>Department of Biochemistry, Yong Loo Lin School of Medicine, National University of Singapore, 117597, Singapore

<sup>§</sup>To whom correspondence should be addressed: pws@caltech.edu



## **Description**

*ina-1* is expressed throughout uterine toroids. At [mid L4](#), expression is seen in uterine toroid 1 and 4 ([ut1](#), [ut4](#)) and in the anterior uterine toroids 2 and 3 ([ut2](#), [ut3](#)). In addition, *ina-1*::GFP is detected in the [spermathecal-uterine junction](#) and [spermatheca](#). At the [L4 lethargus](#) stage, bright expression is observed in uterine toroids 1 and 4, the [spermathecal-uterine junction](#), and [spermatheca](#). Dimmer expression is present in [uterine toroid 2](#) and [posterior toroid 3](#). Fluorescence is also detected in [vulE](#).

## **Reagents**

*gmIs5*

**Funding:** National Institutes of Health USPHS training grant

**Reviewed By:** [Bhagwati Gupta](#)

**History:** Received July 1, 2015 Accepted August 1, 2015 Published March 1, 2016

**Copyright:** © 2016 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Citation:** Ghosh, S; Inoue, T; Sternberg, P (2016). *ina-1* expression in the somatic gonad during late larval development. microPublication Biology. <https://doi.org/10.17912/W2159J>