




A quantitative trait locus for nictation behavior on chromosome V

Jun Kim¹, Daehan Lee^{1,2} and Junho Lee^{1§}

¹Department of Biological Sciences, Institute of Molecular Biology and Genetics, Seoul National University, Seoul 08826, Korea

²Department of Molecular Biosciences, Northwestern University, Evanston, IL, 60208, USA

§To whom correspondence should be addressed: elegans@snu.ac.kr



Nictation ratio of [N2](#) x [CB4856](#) introgression lines on chromosome V. Note that the genotypes of some of these introgression lines including *yslR50* and *yslR51* are estimates from polymorphism markers, not from whole-genome sequencing. Statistics were analyzed by GraphPad Prism and its One-way ANOVA and Dunnett post-test. All tested groups passed the D'Agostino and Pearson omnibus normality test. Each bar indicates the median of each group. *** $P < 0.001$, significantly different from the control strain, N2.



Description

Lee et al., 2017 showed that [CB4856](#) has a lower nictation ratio (the number of nictating dauers over the number of moving dauers on a micro-dirt chip) than [N2](#), and that a genetic locus (*nict-1*) on chromosome IV mediates this phenotype difference. Although this paper had no evidence of any other genetic loci for nictation behavior, we have studied other quantitative trait loci (QTL) using different linkage-based mapping strategies. Here, we report another genetic locus that was identified using introgression lines on chromosome V.

[N2](#) x [CB4856](#) introgression lines were tested using a population nictation assay (Lee et al., 2015). We measured the nictation ratios of the *ewIR65*, *ewIR67*, *ewIR68*, and *ewIR70* lines, which contain N2-type genome in all chromosomes except for [CB4856](#)-type chromosome V segments (Figure 1A). Among them, only *ewIR70* line showed significant lower nictation ratio. Therefore, we narrowed down the chromosome V right region.

We also measured nictation ratios of *ysIR50*, *ewIR74*, and *ysIR51* lines (Figure 1B). *ysIR50* and *ysIR51* were derived from [N2](#) x *ewIR70*. *ysIR50*, *ewIR74*, and *ysIR51* lines showed significantly lower nictation ratios than that of [N2](#). Because the putatively responsible QTL region is covered by another near-isogenic region, *eanIR158*, which was made independently by Erik Andersen's laboratory, we measured the nictation ratios of *eanIR158* and *gyIR3* lines and found that they did not show any significant difference from N2 (Figure 1C).

eanIR158 and *gyIR3* lines were built with [QX1430](#) x N2 (personal communication) and [QX217](#) x N2 (Balla et al., 2015), respectively, and [QX1430](#) and [QX217](#) were made by Leonid Kruglyak's laboratory (Andersen et al., 2015; Rockmen and Kruglyak, 2009). However, *ewIR70* and *ewIR74* were built by Jan Kammenga's laboratory (Doroszuk et al., 2009). Thus, we concluded that the two laboratories may have used different [N2](#) or [CB4856](#) strain derivatives, which may have altered genetic variations in the introgression lines. A take-home lesson is that it is important to examine NILs derived from common original strains in order to follow QTL without being affected by differences in background genotypes.

Jan Kammenga kindly provided their N2 x [CB4856](#) introgression lines for chromosome V, which include *ewIR65*, *ewIR67*, *ewIR68*, *ewIR70*, and *ewIR74*, and Erik Andersen also gave us many introgression lines, which include *eanIR158* and *gyIR3* covering 15-19 Mb of chromosome V. We appreciate their help.

Reagents

Strains: [N2](#), [LJ1250](#) *ysIR50* (V, [CB4856](#)>[N2](#)), [LJ1251](#) *ysIR51* (V, [CB4856](#)>[N2](#)), [ECA238](#) *eanIR158* (V, [CB4856](#), [N2](#)), and [ERT248](#) *gyIR3* (V, [CB4856](#)>[N2](#)).

Genotypes: *ewIR65* (V, [CB4856](#)>[N2](#)), *ewIR67* (V, [CB4856](#)>[N2](#)), *ewIR68* (V, [CB4856](#)>[N2](#)), *ewIR70* (V, [CB4856](#)>[N2](#)), and *ewIR74* (V, [CB4856](#)>[N2](#)).

References

- Andersen EC, Shimko TC, Crissman JR, Ghosh R, Bloom JS, Seidel HS, et al. A Powerful New Quantitative Genetics Platform, Combining *Caenorhabditis elegans* High-Throughput Fitness Assays with a Large Collection of Recombinant Strains. *G3 (Bethesda)*. 2015 Mar 13;5(5):911-20. DOI: 10.1534/g3.115.017178 | PMID: 25770127. | PMCID: PMC4426375.
- Balla KM, Andersen EC, Kruglyak L, Troemel ER. A Wild C. *Elegans* Strain Has Enhanced Epithelial Immunity to a Natural Microsporidian Parasite. *PLOS Pathog*. 2015 Feb 13;11(2):e1004583. DOI: 10.1371/journal.ppat.1004583 | PMID: 25680197. | PMCID: PMC4334554.
- Doroszuk A, Snoek LB, Fradin E, Riksen J, Kammenga J. A genome-wide library of CB4856/N2 introgression lines of *Caenorhabditis elegans*. *Nucleic Acids Res*. 2009 Sep;37(16):e110. DOI: 10.1093/nar/gkp528 | PMID: 19542186. | PMCID: PMC2760803.
- Lee D, Lee H, Choi M-k, Park S, Lee J. Nictation Assays for *Caenorhabditis* and Other Nematodes. *Bio Protoc*. 2015 May 4;5(7):e1433. DOI: 10.1038/nn.2975 | PMID: 22081161.
- Lee D, Yang H, Kim J, Brady S, Zdraljevic S, Zamanian M, et al. The genetic basis of natural variation in a phoretic behavior. *Nat Commun*. 2017;8(1):273 PubMed PMID: 28819099. DOI: 10.1038/s41467-017-00386-x | PMID: 28819099. | PMCID: PMC5561207.
- Rockman MV, Kruglyak L. Recombinational Landscape and Population Genomics of *Caenorhabditis elegans*. *PLoS Genet*. 2009 Mar;5(3):e1000419 DOI: PMC2652117 | PMID: 19283065. | PMCID: 10.1371/journal.pgen.1000419.



9/19/2017 - Open Access

Funding: This work was supported by the National Research Foundation of Korea (NRF) grants funded by the Korean government (MSIT) (No. 2016K1A3A1A47921615). Jun Kim was supported by POSCO Science Fellowship from the POSCO TJ Park Foundation.

Reviewed By: [Erik Andersen](#)

History: **Received** August 24, 2017 **Accepted** September 13, 2017 **Published** September 19, 2017

Copyright: © 2017 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Kim, J; Lee, D; Lee, J (2017). A quantitative trait locus for nictation behavior on chromosome V. *microPublication Biology*. <https://doi.org/10.17912/W23D39>