

lin-41 controls dauer formation and morphology via lin-29 in C. elegans

Allison R Cale^{1,2} and Xantha Karp^{3§}

¹Department of Chemistry and Biochemistry, Central Michigan University, Mount Pleasant, MI 48859

²Current address: Department of Human Genetics, University of Michigan, Ann Arbor, MI 48109

³Department of Biology, Central Michigan University, Mount Pleasant, MI 48859

[§]To whom correspondence should be addressed: karp1x@cmich.edu



Figure 1: *lin-41(0); daf-7* mutants display a *lin-29*-dependent dauer defective phenotype: (A) The *daf-7(e1372)* allele was used to promote dauer entry, and was present in all strains. Larvae homozygous for the *lin-41* null allele *n2914* ("*lin-41(0)*") rarely entered dauer at the dauer-inducing temperature of 24°C, in comparison to control larvae that were wild-type for *lin-41* or heterozygous for *lin-41(0)*. *lin-41(0)* was balanced with the closely-linked transgene *nIs408[LIN-29::mCherry, ttx-3::gfp]* (Harris and Horvitz 2011). *lin-41(0)* homozygous larvae were identified by lack of fluorescence, whereas *lin-41(0)/+* or + refers to larvae that were positive for fluorescence. Simultaneous loss of *lin-29 ("lin-29(0)")* suppressed the dauer-defective

phenotype of *lin-41(0)*, indicating that misregulation of *lin-29* inhibits dauer entry in *lin-41(0)* larvae. (B) SDS-selection from starved plates yielded *lin-41(0)* homozygous dauer larvae, over half of which displayed morphological defects. Morphological defects were absent in *lin-41(0)*; *lin-29(0)* dauers from starved plates, indicating that misregulation of *lin-29* can result in abnormalities in dauer morphology. (C) *daf-7* dauer larvae (top) from starved plates displayed distinct dauer alae with crisp, parallel lines as well as the long, thin body shape characteristic of dauer larvae. By contrast, some *lin-41(0)* dauer larvae displayed morphological defects including crooked and convergent alae and a lumpy body shape (bottom). Images were taken using a 63X objective. (D) Control *lin-41(0)/nIs408* or *nIs408* dauer larvae were long and thin, whereas some *lin-41(0)* homozygous dauer larvae appeared Dpy. Images were taken using a 10X objective. Fisher exact test: ns, not significant; ****, P<0.0001.

Description

In response to adverse environments *C. elegans* larvae may form stress-resistant and developmentally arrested dauer larvae. Entry into dauer interrupts developmental progression after the second larval molt (Cassada and Russell 1975), and the dauer formation decision must be coordinated with developmental pathways. Heterochronic genes control stage-specific cell fate decisions, and these genes interconnect with the dauer formation (*daf*) pathway in several ways. Dauer formation and *daf* genes affect heterochronic phenotypes, and conversely, heterochronic genes can impact dauer by controlling the timing of dauer formation, the morphology of dauer larvae, or the ability to enter dauer (Liu and Ambros 1989; Liu and Ambros 1991; Antebi *et al.* 1998; Tennessen *et al.* 2010; Karp and Ambros 2011; Karp and Ambros 2012). Here we find that the *lin-41* heterochronic gene regulates the dauer formation decision and dauer morphogenesis via its canonical target, *lin-29*.

The temperature-sensitive daf-7(e1372) allele results in constituitive dauer entry at 25°C, and the allele is a useful tool for controlling dauer entry (Vowels and Thomas 1992; Karp 2018). While performing experiments examining the role of *lin-41* during dauer, we observed that *lin-41(0); daf*-7 worms rarely entered dauer. To better understand this phenotype, we quantified the number of dauer and nondauer larvae in synchronous populations of *daf*-7(e1372) larvae grown at the dauer-inducing temperature of 24°C.

lin-41(0) mutants are sterile, and therefore the strain must be propagated as a heterozygote (Slack *et al.* 2000). We took advantage of a closely linked transgene, *nIs408[LIN-29::mCherry, ttx-3::gfp]* to balance *lin-41(0)* (Harris and Horvitz 2011). *lin-41(0)* homozygous larvae were identified by the absence of fluorescence. Controls included *nIs408*-positive siblings, 2/3 of which should be *lin-41(0)/nIs408* and 1/3 of which should be *nIs408/nIs408*, as well as a strain that is homozygous for the wild-type allele of *lin-41*.

Embryos from each strain were incubated at 24°C for 50-52 hours to induce dauer formation. As expected, 100% of larvae in the control *daf-7* strain were dauers (Fig. 1A). Similarly, the *lin-41(0)/nIs408* or *nIs408/nIs408; daf-7* larvae formed dauers at high penetrance (96%). By contrast, only 2/23 (9%) of *lin-41(0)* homozygous larvae were in dauer. The remaining *lin-41(0)* larvae had continued development and were in the L3 or L4 stage (Fig. 1A). Therefore, *lin-41(0)* larvae display a dauer formation-defective (Daf-d) phenotype.

The canonical role for *lin-41* as a heterochronic gene is to promote larval cell fate in hypodermal cells by directly repressing translation of *lin-29* (Slack *et al.* 2000; Aeschimann *et al.* 2017). *lin-29* encodes a transcription factor that promotes adult cell fate (Rougvie and Ambros 1995). In addition to this canonical role, *lin-41* controls several other processes, including oocyte growth and maturation, male tail tip morphogenesis, and a sex-specific neurotransmitter switch in the AIM interneurons, among others. *lin-41* acts via *lin-29* to control some of these processes, and via other targets to control other processes (Del Rio-Albrechtsen *et al.* 2006; Spike *et al.* 2014; Pereira *et al.* 2019).

To ask whether *lin-29* is required for the Daf-d phenotype we observed in *lin-41(0)* mutants, we examined *lin-41(0); lin-29(0)* larvae segregating from *lin-41(0)/nIs408; lin-29(0)* mothers. Because the *nIs408* transgene rescues *lin-29(0)* (Harris and Horvitz 2011), this transgene effectively balances both *lin-41* and *lin-29*. We found that the additional loss of *lin-29(0)* suppressed the *lin-41(0)* dauer entry-defective phenotype, as 96% of *lin-41(0); lin-29(0); daf-7* larvae examined were in dauer (Fig. 1A).

Since obtaining *lin-41(0)* dauers using the *daf-7* method was inefficient, we asked whether utilization of the natural starvation response could drive *lin-41(0)*; *daf-7* larvae into dauer. We allowed each strain to crowd and starve out, and then selected for dauer larvae using 1% SDS (Cassada and Russell 1975). We were able to obtain SDS-resistant *lin-41(0)* homozygous dauer larvae using this method. Forty-two percent of *lin-41(0)* dauer larvae displayed normal dauer morphology, as did the two dauer larvae obtained from *daf-7* induced dauer formation (Fig. 1A). Interestingly, the remaining 58% of *lin-41(0)* dauer larvae displayed defective dauer morphology (Fig. 1B). Specifically, both dauer alae and body shape were abnormal. In contrast to the straight ridges that make up wild-type dauer alae, *lin-41(0)* mutant dauers displayed wavy dauer alae (Fig. 1C). With

respect to body shape, there was a range in severity from worms of a normal length but a somewhat lumpy appearance to Dpy dauers that were shorter and fatter than controls (Fig. 1D).

These dauer morphology defects are reminiscent of those caused by loss of either of two other heterochronic genes: *lin-14* or *lin-28* (Liu and Ambros 1989). However, those defects appear to be due to areas of lateral cuticle that lack dauer alae. Interestingly, the *lin-28(0)* defect is suppressed by loss of *lin-29*, whereas the *lin-14* defect is not (Liu and Ambros 1989). By contrast, we never observed gaps in dauer alae in *lin-41(0)* mutants. Furthermore, similarly to the Daf-d phenotype, we found that the dauer morphology phenotype observed in the *lin-41(0)* homozygotes was suppressed by loss of *lin-29*, as no *lin-41(0)*; *lin-29(0)*; *daf-7* dauer larvae from the SDS experiments displayed defective dauer morphology (Fig. 1B).

In summary, *lin-41* regulates two aspects of dauer biology, the decision to enter dauer, and dauer morphogenesis. In both cases, *lin-41* appears to act via its canonical target, *lin-29*. This finding adds to our understanding of the apparent coordination between dauer formation pathways and heterochronic pathways.

Methods

Request a detailed protocol

Strains and maintenance.

C. elegans strains were maintained at 20°C on NGM plates with *E. coli* strain OP50 as a food source (Brenner 1974).

Dauer-induction via daf-7 and dauer formation assay

Individual gravid adult hermaphrodites from each strain were transferred to 60mm NGM plates and allowed to lay embryos for 3-5 hours at 24°C. Adults were then removed, and embryos were incubated at 24°C for an additional 50 to 52 hours. For heterozygous strains, any plates containing homozygous mothers were discarded. Larvae were then imaged as described below. For strains XV238 (*lin-41/nIs408; daf-7*) and the control VT1777 (*daf-7*), every larva on the plate was scored for developmental stage. For strain XV239 *lin-41/nIs408; lin-29; daf-7*), only worms that lacked *nIs408* were scored, since *nIs408* balances *lin-41* and rescues *lin-29*. Non-dauer larvae were distinguishable from dauer larvae primarily due to the lack of dauer alae, in addition to their larger size, lack of radial constriction, and progression of gonad and vulval development. Six independent trials were performed.

Dauer induction via starvation and dauer morphology assay

Strains were propagated at 20°C on 60mm NGM plates seeded with OP50. Heterozygous strains were maintained by cloning out individuals and verifying the correct segregation of phenotypes in the progeny. Plates were monitored to identify the point at which the food was exhausted and the point at which dauer larvae appeared. Sets of XV238, XV239, and VT1777 populations that starved at the same time were paired for experiments, to ensure that dauer larvae were approximately the same age between strains within a single experiment. The data in Figure 1B represent 6 independent trials. Dauer larvae were selected using a 30-35 minute incubation in 1% SDS at room temperature. For strain XV239, only larvae lacking fluorescence were analyzed.

Microscopy

Larvae were picked to slides with 2% agarose pads and anesthetized in 0.1 M levamisole. Worms were visualized using a Zeiss AxioImager D2 compound microscope with DIC optics. Representative images acquired using an AxioCam MRm Rev 3 camera and Zen software (2012) from Zeiss are shown. Brightness/contrast was adjusted using Adobe Photoshop.

Reagents

Strains used in this study

Strain name	Genotype
VT1777	daf-7(e1372); maIs105[col-19::gfp]
XV238	lin-41(n2914)/nIs408[LIN-29::mCherry, ttx-3::gfp]; daf-7(e1372); maIs105
XV239	lin-41(n2914)/nIs408; lin-29(n546); daf-7(e1372); maIs105

Acknowledgments: Thanks to Matthew Wirick (Karp lab) for help with some of the imaging, Anna Zinovyeva (Kansas State University), and Himal Roka (Karp lab) for critical reading of the manuscript, and to WormBase for information. Some strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440).



References

Aeschimann F, Kumari P, Bartake H, Gaidatzis D, Xu L, Ciosk R, Großhans H. 2017. LIN41 Post-transcriptionally Silences mRNAs by Two Distinct and Position-Dependent Mechanisms. Mol Cell 65: 476-489.e4. PMID: 28111013.

Antebi A, Culotti JG, Hedgecock EM. 1998. *daf-12* regulates developmental age and the dauer alternative in *Caenorhabditis elegans*. Development 125: 1191-205. PMID: 9477318.

Brenner S. 1974. The genetics of Caenorhabditis elegans. Genetics 77: 71-94. PMID: 4366476.

Cassada RC, Russell RL. 1975. The dauerlarva, a post-embryonic developmental variant of the nematode *Caenorhabditis elegans*. Dev Biol 46: 326-42. PMID: 1183723.

Del Rio-Albrechtsen T, Kiontke K, Chiou SY, Fitch DH. 2006. Novel gain-of-function alleles demonstrate a role for the heterochronic gene *lin-41* in *C. elegans* male tail tip morphogenesis. Dev Biol 297: 74-86. PMID: 16806150.

Harris DT, Horvitz HR. 2011. MAB-10/NAB acts with LIN-29/EGR to regulate terminal differentiation and the transition from larva to adult in *C. elegans*. Development 138: 4051-62. PMID: 21862562.

Karp X. 2018. Working with dauer larvae. WormBook 2018: 1-19. PMID: 27417559.

Karp X, Ambros V. 2011. The developmental timing regulator HBL-1 modulates the dauer formation decision in *Caenorhabditis elegans*. Genetics 187: 345-53. PMID: 20980238.

Karp X, Ambros V. 2012. Dauer larva quiescence alters the circuitry of microRNA pathways regulating cell fate progression in *C. elegans*. Development 139: 2177-86. PMID: 22619389.

Liu Z, Ambros V. 1991. Alternative temporal control systems for hypodermal cell differentiation in *Caenorhabditis elegans*. Nature 350: 162-5. PMID: 26502479.

Liu ZC, Ambros V. 1989. Heterochronic genes control the stage-specific initiation and expression of the dauer larva developmental program in *Caenorhabditis elegans*. Genes Dev 3: 2039-49. PMID: 2628162.

Pereira L, Aeschimann F, Wang C, Lawson H, Serrano-Saiz E, Portman DS, Großhans H, Hobert O. 2019. Timing mechanism of sexually dimorphic nervous system differentiation. Elife 8: . PMID: 30599092.

Rougvie AE, Ambros V. 1995. The heterochronic gene *lin-29* encodes a zinc finger protein that controls a terminal differentiation event in *Caenorhabditis elegans*. Development 121: 2491-500. PMID: 7671813.

Slack FJ, Basson M, Liu Z, Ambros V, Horvitz HR, Ruvkun G. 2000. The *lin-41* RBCC gene acts in the *C. elegans* heterochronic pathway between the *let-7* regulatory RNA and the LIN-29 transcription factor. Mol Cell 5: 659-69. PMID: 10882102.

Spike CA, Coetzee D, Eichten C, Wang X, Hansen D, Greenstein D. 2014. The TRIM-NHL protein LIN-41 and the OMA RNA-binding proteins antagonistically control the prophase-to-metaphase transition and growth of *Caenorhabditis elegans* oocytes. Genetics 198: 1535-58. PMID: 25261698.

Tennessen JM, Opperman KJ, Rougvie AE. 2010. The *C. elegans* developmental timing protein LIN-42 regulates diapause in response to environmental cues. Development 137: 3501-11. PMID: 20843862.

Vowels JJ, Thomas JH. 1992. Genetic analysis of chemosensory control of dauer formation in *Caenorhabditis elegans*. Genetics 130: 105-23. PMID: 1732156.

Funding: NIH R15GM117568 and NSF IOS CAREER 1652283

Author Contributions: Allison R Cale: Conceptualization, Formal analysis, Investigation, Methodology, Visualization, Writing - original draft, Writing - review and editing. Xantha Karp: Conceptualization, Funding acquisition, Methodology, Project administration, Supervision, Writing - review and editing.

Reviewed By: Anonymous

History: Received October 28, 2020 Revision received November 5, 2020 Accepted November 9, 2020 Published November 12, 2020

Copyright: © 2020 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Citation: Cale, AR; Karp, X (2020). *lin-41* controls dauer formation and morphology via *lin-29* in *C. elegans*. microPublication Biology. https://doi.org/10.17912/micropub.biology.000323