

Field Sampling of *Zaprionus indianus* in the Eastern United States in 2023-2024

Weston J. Gray¹, Megan Stephenson², Alan O. Bergland², Priscilla A. Erickson^{1§}

¹Biology, University of Richmond, Richmond, VA, USA

²Biology, University of Virginia, Charlottesville, VA, USA

[§]To whom correspondence should be addressed: perickso@richmond.edu

Abstract

We conducted surveys for the invasive drosophilid *Zaprionus indianus* in the eastern United States in 2023 and 2024. We found no latitudinal trends in *Z. indianus* abundance, and the northern boundary for *Z. indianus* was variable between years. *Z. indianus* was rare in central and northern Florida in the spring, was not present on early season berry crops in Virginia, and had a restricted temperate growing season compared to other drosophilids, providing further support for the species' yearly recolonization of temperate areas. *Z. indianus* was also observed on native fruits in a natural area in Virginia.




Figure 1. *Z. indianus* survey results from 2023 and 2024:

All data are shown as the proportion of *Z. indianus* relative to other drosophilids from collections in 2023 and 2024. A-B) Results of latitudinal sampling from 2023 (A) and 2024 (B). C) No *Z. indianus* were collected at two orchards in Massachusetts in either year. D) *Z. indianus* relative abundance varied between sites in southern Florida in November 2023. E-F) Collections conducted in Florida and southern Georgia in March 2024 (E) and May 2024 (F). G-H) Virginia sampling results during the 2023 (G) and 2024 (H) growing seasons. Point size is scaled to the total number of drosophilids captured. Open circles indicate Carter Mountain (CM) collections using nets and aspirators, while the other CM collections (closed circles) were conducted via trapping. HPO=Hanover Peach Orchard; ABF = Agriberry Farm. I) Incidental observation of *Z. indianus* on fallen fruit of *Asimina triloba* (pawpaw, top) and drupes of *Juglans nigra* (black walnut, bottom) in a natural environment in Richmond, VA. *Z. indianus* are circled; arrowhead points to a copulating pair.

Description

One of the most biologically damaging consequences of globalization is the introduction and spread of invasive species to foreign environments, which can cause irreparable damage to natural ecosystems and inflict severe economic costs (Mooney & Cleland, 2001; Holmes et al., 2009; Oliveira et al., 2013). *Zaprionus indianus*, the African fig fly, is an invasive drosophilid native to tropical Africa that has since spread to India, the Middle East, both North and South

America, and to a limited extent Europe (Vilela, 1999; Yassin et al., 2008; Al-Jboory & Katbeh-Bader, 2012; EFSA Panel on Plant Health (PLH) et al., 2022; Nagy et al., 2025). In the western hemisphere, *Z. indianus* was first reported in Brazil in 1999 and subsequently spread across the continent and into North America (Vilela, 1999; Goñi et al., 2001; Van der Linde et al., 2006; Markow et al., 2014). *Z. indianus* was first identified in Florida in 2005 (Van der Linde et al., 2006), was reported in Virginia in 2012 (Pfeiffer et al., 2019), and has since been detected as far north as Minnesota, Quebec, and Ontario (Renkema et al., 2013; Holle et al., 2018).

Z. indianus often dominates local drosophilid communities, especially during warmer months, and negatively impacts their diversity (Tidon et al., 2003; Silva et al., 2005; Roque et al., 2017; Pfeiffer et al., 2019; Rakes et al., 2023). Furthermore, *Z. indianus* is an agricultural pest and poses a potential economic threat for fig and berry production, as it is capable of ovipositing on healthy fruits prior to harvest, leading to larval damage of the ripening fruit (Stein et al., 2003; Oliveira et al., 2013; Bernardi et al., 2017; Allori Stazonelli et al., 2023; EFSA Panel on Plant Health (PLH) et al., 2022). As such, efforts to track the dynamics of its ongoing invasion play an important role in developing effective management strategies and understanding the intricacies of its invasion biology.

We surveyed *Z. indianus* in the eastern United States along a latitudinal transect spanning Florida to Massachusetts in the fall of 2023 and 2024. *Z. indianus* was generally common and widespread; the species comprised over 50% of the total drosophilids collected at many sites (Figure 1A, 1B, & 1D). Our findings show high variability between localities, as surveys conducted at nearby locations on the same day produced considerably different proportions of *Z. indianus* (Figure 1D, Rakes et al. 2023). However, some of this variability is likely a result of conducting adjacent sampling efforts at different times of day, since fly activity may vary due to differences in temperature or circadian rhythms. Unlike our sampling from 2022 (Rakes et al., 2023), we did not detect *Z. indianus* in Massachusetts in either 2023 or 2024 (Figure 1C), and *Z. indianus* was rare in Connecticut in 2023. However, in 2024, we detected *Z. indianus* in Albany, New York, and their abundance was substantially higher in Connecticut, indicating a reduced range in 2023 relative to 2022 and 2024. Taken together, these findings suggest the northern boundary of its invasion fluctuates yearly. This observation is supported by other studies that have demonstrated similar patterns of annual variability in abundance at higher latitudes (Holle et al., 2018; Dettler et al., 2021; Nagy et al., 2025). Lastly, there was no correlation between abundance and latitude in either 2023 or 2024 (Pearson correlation, $t = -0.697$, $df = 7$, $P = 0.508$, $r = -0.255$ and $t = 0.135$, $df = 5$, $P = 0.898$, $r = 0.06$, respectively).

We also conducted temporal sampling of *Z. indianus* at two peach and apple orchards in Virginia in 2023 and 2024. Similar to prior surveys (Rakes et al., 2023), *Z. indianus* did not appear each year until late in the summer, at which point the population expanded quickly within a month and usually became the dominant drosophilid, before slowly decreasing and disappearing before the onset of winter in late November/early December (Figure 1G & 1H). Sampling method also influenced collections; we often captured a greater proportion of *Z. indianus* by netting and aspirating than using traps at Carter Mountain (Figure 1G & 1H, compare open and closed circles).

Our findings of a delayed emergence and early disappearance compared to other drosophilids suggest that *Z. indianus* currently undergoes an annual northward recolonization event from southern refugia before being extirpated in the winter (Pfeiffer et al., 2019; Rakes et al., 2023). While we only conducted detailed temporal sampling at a single latitude, studies conducted at localities north of Virginia did not report *Z. indianus* until later in the season than our first detections, suggesting a stepwise progression northwards throughout the growing season (Joshi et al., 2014; Holle et al., 2018). We speculate that this northward expansion is primarily mediated by commercial produce transport as fruits such as peaches and berries ripen earlier at southern latitudes and are shipped north, though we cannot rule out the possibility of natural causes such as wind or migratory flight. Alternatively, *Z. indianus* could overwinter and emerge at different times at different latitudes, but our data and other studies suggest overwintering does not occur in temperate populations. We found few *Z. indianus* in central and northern Florida in the spring, suggesting even mild winters limit population sizes (Figure 1E & 1F). Another study showed limited populations of *Z. indianus* in northern Florida during the winter and spring, while populations remained abundant in central/southern Florida (Renkema et al., 2018). Likewise, no *Z. indianus* were collected during winter sampling efforts in Memphis, Tennessee (Kohlmeier & Kohlmeier, 2025). Furthermore, although a handful of studies have demonstrated some cold-tolerant traits in *Z. indianus* (Ramniwas et al., 2012; Lavagnino et al., 2020), many others have demonstrated limitations in freezing conditions (Karan et al., 1999; Araripe et al., 2004; Nava et al., 2007; Comeault et al., 2020; Lavagnino et al., 2020). While our findings are consistent with annual recolonization, additional fine-grained temporal sampling and physiological studies are needed to definitively reject the hypothesis of overwintering.

We also monitored a Virginia berry farm in 2024 to determine whether *Z. indianus* were present on berries in the early summer, prior to peaches ripening at other locales later. No *Z. indianus* were detected at the berry farm prior to other locations in Virginia, and they remained uncommon throughout the fall even while abundant at surrounding orchards (Figure 1H, gray). Previous studies have reported mixed results regarding *Z. indianus* abundances near brambleberries (Joshi et al., 2014; Holle et al., 2018; Dettler et al., 2021; Allori Stazonelli et al., 2023), and differential use of pesticides could explain variation in abundances between sites (Andreazza et al., 2016; de Oliveira Rios et al., 2024). While our results collectively suggest *Z. indianus* arrives in Virginia later in the summer, the lack of *Z. indianus* on berries earlier in

the season may instead be a product of the local environment at the berry farm we sampled, rather than a true reflection of the species' absence in Virginia at that time. Our finding that *Z. indianus* emerges in Virginia at approximately the same time each year demonstrates relative consistency in the timing of expansion. We suggest that this timing pattern is largely due to the structured seasonality of produce production and transport. However, *Z. indianus* has previously been collected in Virginia as early as June, suggesting additional variables impact the timing of its annual expansion, or that overwintering could occur in some years (Erickson et al., 2025).

Apart from our planned collections, we incidentally found evidence of *Z. indianus* aggregating and copulating on fallen pawpaw (*Asimina triloba*) fruits and drupes of the black walnut (*Juglans nigra*) in a natural park in Richmond, Virginia (Figure 1I). To our knowledge, this is the first report of *Z. indianus* utilizing either species as a host (EPPO, 2025). Further examination will be required to determine whether these native fruits are suitable larval hosts. Previous studies have found *Z. indianus* utilizing native fruit to varying success, suggesting the species is invading both agricultural and native ecosystems (Leão & Tidon, 2004; Castrenza, 2011; Joshi et al., 2014). The lack of commercial produce production or transport at this location suggests that short-range dispersal of *Z. indianus* can occur independently of fruit shipments. Thus, the use of wild hosts in natural areas could permit further range expansion without human assistance and would alter the approach to formulating effective management strategies. Furthermore, this expansion to natural habitats could pose additional threats to native drosophilids, which often find refuge from invasive species in these areas (Tidon et al., 2003; Hochmüller et al., 2010; Liu et al., 2023).

Z. indianus can undergo rapid evolution in its invaded range (Comeault et al., 2020, 2021; Erickson et al., 2025; Gray et al., 2025). In the future, the species could potentially evolve cold tolerance adaptations that would allow successful overwintering in temperate areas. Year-round populations could have adverse effects on both the local fly communities and agricultural industry of these regions without the limiting factor of annual extirpation. As such, ongoing monitoring is vital to develop and implement effective strategies to limit impacts on both natural and agricultural systems and to track changes that could worsen its invasive potential.

Methods

Two orchards in Virginia, Carter Mountain Orchard (37.992, -78.472) and Hanover Peach Orchard (37.570, -77.265), were utilized for temporal collections in 2023 and 2024. Both orchards produced peaches in the summer and apples in the fall, and sampling efforts centered around the fruit in season. We collected approximately every 2-4 weeks from June through late November/early December to track the seasonal fluctuations of *Z. indianus*. In 2024, we sampled at Agriberry Farm (37.726, -77.314) to assess the fruit fly community in Virginia during the early summer on blackberries and raspberries. Latitudinal sampling from Florida to Massachusetts took place each fall in both 2023 and 2024, with most collections conducted in October (see Table for locations and dates). Additional collections were conducted in the spring of 2024 in Florida (primarily citrus) and southern Georgia (blueberries) to estimate the northern limits of the year-round population of *Z. indianus* and the dynamics of its initial range expansion north each year.

We netted over fallen fruit and conducted unbiased aspiration directly off the fruit on the ground. Sampling method choice was dependent on a combination of environmental factors such as temperature, precipitation, and abundance of fallen fruit. Traps were used for all Carter Mountain collections (unless otherwise noted) and as needed at some locations. Traps were constructed from 2 L plastic bottles baited with sliced bananas sprinkled with baker's yeast and were hung from trees or trellises overnight. Flies were collected from the traps the next day via aspiration. See Rakes et al. 2023 for additional sampling methodology details. Collected flies were anaesthetized with CO₂ and sorted under a stereomicroscope to determine counts of *Z. indianus* and other drosophilids. *Z. indianus* were photographed with an iPhone on pawpaws and black walnuts at Pony Pasture Park (37.551, -77.517) in Richmond, Virginia on September 19, 2024. A subsequent flood prevented us from returning to collect flies.

Data were processed using *data.table* (Barrett et al., 2024) in R (R Core Team, 2024, v.4.3.3) and figures were created using *ggplot2* (Wickham, 2016) and *cowplot* (Wilke, 2024). We conducted Pearson's correlation analyses to test for a relationship between latitude and the proportion of *Z. indianus* collected.

Reagents

| State | Orchard | Date of Collection | Latitude | Longitude | Relative Abundance | Total Collected |
|-------|-------------------|--------------------|----------|-----------|--------------------|-----------------|
| NY | Bowman Orchards | 10/13/2024 | 42.817 | -73.839 | 0.08 | 110 |
| MA | Westward Orchards | 10/14/2024 | 42.494 | -71.561 | 0 | 198 |

| | | | | | | |
|----|-------------------------------------|------------|--------|---------|------|------|
| MA | Westward Orchards | 10/10/2023 | 42.494 | -71.561 | 0 | 241 |
| MA | Carver Hill Orchard | 10/14/2024 | 42.430 | -71.504 | 0 | 203 |
| MA | Carver Hill Orchard | 10/10/2023 | 42.430 | -71.504 | 0 | 200 |
| CT | Lyman Orchards | 10/15/2024 | 41.494 | -72.730 | 0.67 | 273 |
| CT | Lyman Orchards | 10/11/2023 | 41.494 | -72.730 | 0.02 | 450 |
| PA | Linvilla Orchards | 10/16/2024 | 39.885 | -75.410 | 0.91 | 309 |
| PA | Linvilla Orchards | 10/12/2023 | 39.885 | -75.410 | 0.6 | 636 |
| NC | Sky Top Orchard | 9/24/2023 | 35.240 | -82.436 | 0.7 | 44 |
| GA | Hillcrest Orchards | 10/11/2024 | 34.620 | -84.373 | 0.47 | 245 |
| GA | Hillcrest Orchards | 9/25/2023 | 34.620 | -84.373 | 0.16 | 1271 |
| GA | Oglesby Family Farm | 5/7/2024 | 31.144 | -81.621 | 0.02 | 65 |
| GA | Morning Belle Farms | 5/8/2024 | 30.864 | -81.635 | 0.03 | 142 |
| FL | Kent's Blueberry Farm | 5/8/2024 | 30.488 | -81.738 | 0 | 29 |
| FL | Showcase of Citrus | 3/29/2024 | 28.424 | -81.704 | 0 | 88 |
| FL | Schacht Groves | 3/29/2024 | 27.626 | -80.452 | 0.2 | 673 |
| FL | University of Miami | 11/11/2023 | 25.722 | -80.280 | 0.01 | 529 |
| FL | The Fairchild Farm | 11/11/2023 | 25.537 | -80.433 | 0.28 | 774 |
| FL | Fruit & Spice Park | 10/22/2024 | 25.535 | -80.493 | 0.08 | 1480 |
| FL | Fruit & Spice Park | 3/27/2024 | 25.535 | -80.493 | 0.13 | 512 |
| FL | Schnebly Redland's Winery & Brewery | 11/11/2023 | 25.486 | -80.541 | 0.57 | 67 |

Acknowledgements: We thank the owners and managers of all of the orchards and farms that we visited for permission to collect flies. We thank Camille Walsh-Antzak, Ansleigh Gunter, Dr. Shannon Jones, and the students from the University of Richmond Integrated Science Experience program for assistance with fly collections.

References

- Al Jboory IJ, Katbeh Bader A. 2012. First Record of *Zaprionus indianus* (Gupta, 1970) (Drosophilidae: Diptera) in Jordan. Allori Stazonelli E, Funes CF, Corral Gonzalez MN, Gibilisco SM, Kirschbaum DS. 2023. Population fluctuation and infestation levels of *Zaprionus indianus* Gupta (Diptera: Drosophilidae) in berry crops of northwestern Argentina. Acta Horticulturae: 141. DOI: [10.17660/ActaHortic.2023.1381.19](https://doi.org/10.17660/ActaHortic.2023.1381.19)
- Andreazza F, Bernardi D, Baronio CA, Pasinato J, Nava DE, Botton M. 2016. Toxicities and effects of insecticidal toxic baits to control *Drosophila suzukii* and *Zaprionus indianus* (Diptera: Drosophilidae). Pest Management Science. 73: 146.

DOI: [10.1002/ps.4348](https://doi.org/10.1002/ps.4348)

Araripe LO, Klaczko LB, Moreteau B, David JR. 2004. Male sterility thresholds in a tropical cosmopolitan drosophilid, *Zaprionus indianus*. *Journal of Thermal Biology*. 29: 73. DOI: [10.1016/j.jtherbio.2003.11.006](https://doi.org/10.1016/j.jtherbio.2003.11.006)

Barrett T, Dowle M, Srinivasan A, Gorecki J, Chirico M, Hocking T, Schwendinger B. 2024. data.table: Extension of `data.frame`. R package version 1.16.2. DOI: [10.32614/CRAN.package.data.table](https://doi.org/10.32614/CRAN.package.data.table)

Bernardi D, Andreazza F, Botton M, Baronio CA, Nava DE. 2017. Susceptibility and Interactions of *Drosophila suzukii* and *Zaprionus indianus* (Diptera: Drosophilidae) in Damaging Strawberry. *Neotropical Entomology*. 46: 1. DOI: [10.1007/s13744-016-0423-9](https://doi.org/10.1007/s13744-016-0423-9)

Castrenza S. 2011. New record on novel hosts for the Drosophilid pest *Zaprionus indianus*. *Dros. Inf. Serv.* 94: 1.

Comeault AA, Kautt AF, Matute DR. 2021. Genomic signatures of admixture and selection are shared among populations of *Zaprionus indianus* across the western hemisphere. *Molecular Ecology*. 30: 6193. DOI: [10.1111/mec.16066](https://doi.org/10.1111/mec.16066)

Comeault AA, Wang J, Tittes S, Isbell K, Ingle S, Hurlbert AH, Matute DR. 2020. Genetic Diversity and Thermal Performance in Invasive and Native Populations of African Fig Flies. *Molecular Biology and Evolution*. 37: 1893. DOI: [10.1093/molbev/msaa050](https://doi.org/10.1093/molbev/msaa050)

De Oliveira Rios J, Costa SC, Da Conceicao Galego LG. 2024. Geographical and sex-specific effects of malathion insecticide selection in African fig fly *Zaprionus indianus* Gupta, 1970 (Diptera: Drosophilidae). *Journal of Applied Entomology*. 148: 312. DOI: [10.1111/jen.13212](https://doi.org/10.1111/jen.13212)

Dettler MA, Barrientos GN, Martinez E, Ansa MA, Santadino MV, Coviella CE, Riquelme Virgala MB. 2021. Field infestation level of *Zaprionus indianus* Gupta and *Drosophila suzukii* (Matsumura) (Diptera: Drosophilidae) in *Ficus carica* L. (Rosales: Moraceae) and *Rubus idaeus* L. (Rosales: Rosaceae) in the Northeast of Buenos Aires province. *Revista de la Sociedad Entomológica Argentina*. 80: 43. DOI: [10.25085/rsea.800307](https://doi.org/10.25085/rsea.800307)

EFSA Panel On Plant Health PLH, Bragard C, Baptista P, Chatzivassiliou E, Di Serio F, Gonthier P, et al., MacLeod A. 2022. Pest categorisation of *Zaprionus indianus*. *EFSA Journal*. 20: e07144. DOI: [10.2903/j.efsa.2022.7144](https://doi.org/10.2903/j.efsa.2022.7144)

EPPO EMPPO. 2025. *Zaprionus indianus* (ZAPRIN) Hosts. EPPO Global Database <<https://gd.eppo.int/taxon/ZAPRIN>>

Erickson PA, Stellwagen A, Bangerter A, Gunter A, Polizos NT, Bergland AO. 2025. Limited population structure but signals of recent selection in introduced African Fig Fly (*Zaprionus indianus*) in North America. DOI: [10.1101/2024.09.20.614190](https://doi.org/10.1101/2024.09.20.614190)

Goni B, Fresia P, Calvino M, Ferreiro MJ, Valente LS, Basso Da Silva L. 2001. First record of *Zaprionus indianus* Gupta, 1970 (Diptera, Drosophilidae) in southern localities of Uruguay. *Drosophila Information Service*. 84: 61.

Gray WJ, Rakes LM, Cole C, Gunter A, He G, Morgan S, et al., Erickson PA. 2025. Rapid wing size evolution in African fig flies (*Zaprionus indianus*) following temperate colonization. *Evolution*: qpap102. DOI: [10.1093/evolut/qpap102](https://doi.org/10.1093/evolut/qpap102)

Hochmuller CJC, Lopes Da Silva M, Valente VLS, Schmitz HJ. 2010. The drosophilid fauna (Diptera, Drosophilidae) of the transition between the Pampa and Atlantic Forest Biomes in the state of Rio Grande do Sul, southern Brazil: first records. *Papéis Avulsos de Zoologia*. 50: 286. DOI: [10.1590/S0031-10492010001900001](https://doi.org/10.1590/S0031-10492010001900001)

Holle SG, Tran AK, Burkness EC, Ebbenga DN, Hutchison WD. 2018. First Detections of *Zaprionus indianus* (Diptera: Drosophilidae) in Minnesota. *Journal of Entomological Science*. 54: 99. DOI: [10.18474/JES18-22](https://doi.org/10.18474/JES18-22)

Holmes TP, Aukema JE, Von Holle B, Liebhold A, Sills E. 2009. Economic Impacts of Invasive Species in Forests. *Annals of the New York Academy of Sciences*. 1162: 18. DOI: [10.1111/j.1749-6632.2009.04446.x](https://doi.org/10.1111/j.1749-6632.2009.04446.x)

Joshi NK, Biddinger DJ, Demchak K, Deppen A. 2014. First Report of *Zaprionus indianus* (Diptera: Drosophilidae) in Commercial Fruits and Vegetables in Pennsylvania. *Journal of Insect Science*. 14 DOI: [10.1093/jisesa/ieu121](https://doi.org/10.1093/jisesa/ieu121)

Karan D, Moreteau B, David JR. 1999. Growth temperature and reaction norms of morphometrical traits in a tropical drosophilid: *Zaprionus indianus*. *Heredity*. 83: 398. DOI: [10.1038/sj.hdy.6885940](https://doi.org/10.1038/sj.hdy.6885940)

Kohlmeier P, Kohlmeier P. 2025. A non-overwintering urban population of the African fig fly (Diptera: Drosophilidae) impacts the reproductive output of locally adapted fruit flies. *Florida Entomologist*. 108 DOI: [10.1515/flaent-2024-0064](https://doi.org/10.1515/flaent-2024-0064)

Lavagnino NJ, Fanara JJ, Mensch J. 2020. Comparison of overwintering survival and fertility of *Zaprionus indianus* (Diptera: Drosophilidae) flies from native and invaded ranges. *Journal of Thermal Biology*. 87: 102470. DOI: [10.1016/j.jtherbio.2019.102470](https://doi.org/10.1016/j.jtherbio.2019.102470)

Leao BFD, Tidon R. 2004. Newly invading species exploiting native host-plants: the case of the African *Zaprionus indianus* (Gupta) in the Brazilian Cerrado (Diptera, Drosophilidae). *Annales de la Société entomologique de France (N.S.)*. 40: 285. DOI: [10.1080/00379271.2004.10697427](https://doi.org/10.1080/00379271.2004.10697427)

- Liu D, Semenchuk P, Essl F, Lenzner B, Moser D, Blackburn TM, et al., Dullinger S. 2023. The impact of land use on non-native species incidence and number in local assemblages worldwide. *Nature Communications*. 14: 2090. DOI: [10.1038/s41467-023-37571-0](https://doi.org/10.1038/s41467-023-37571-0)
- Markow TA, Hanna G, Riesgo Escovar JR, Tellez Garcia AA, Richmond MP, Nazario Yepiz NO, et al., Pfeiler E. 2014. Population genetics and recent colonization history of the invasive drosophilid *Zaprionus indianus* in Mexico and Central America. *Biological Invasions*. 16: 2427. DOI: [10.1007/s10530-014-0674-5](https://doi.org/10.1007/s10530-014-0674-5)
- Mooney HA, Cleland EE. 2001. The evolutionary impact of invasive species. *Proceedings of the National Academy of Sciences*. 98: 5446. DOI: [10.1073/pnas.091093398](https://doi.org/10.1073/pnas.091093398)
- Nagy C, Sen E, Kiss B. 2025. First record of African fig fly, *Zaprionus indianus* Gupta, 1970 (Diptera, Drosophilidae) in Hungary. *Bulletin of Insectology*. 78: 27. DOI: [10.3897/bull.insectology.154143](https://doi.org/10.3897/bull.insectology.154143)
- Nava DE, Nascimento AM, Stein CP, Haddad ML, Bento JMS, Parra JRP. 2007. Biology, thermal requirements, and estimation of the number of generations of *Zaprionus indianus* (Diptera: Drosophilidae) for the main fig producing regions of Brazil. *Florida Entomologist*. 90: 495. DOI: [10.1653/0015-4040\(2007\)90\[495:BTAEQ\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2007)90[495:BTAEQ]2.0.CO;2)
- Oliveira CM, Auad AM, Mendes SM, Frizzas MR. 2013. Economic impact of exotic insect pests in Brazilian agriculture. *Journal of Applied Entomology*. 137: 1. DOI: [10.1111/jen.12018](https://doi.org/10.1111/jen.12018)
- Pfeiffer DG, Shrader ME, Wahls JCE, Willbrand BN, Sandum I, Van Der Linde K, et al., Day ER. 2019. African Fig Fly (Diptera: Drosophilidae): Biology, Expansion of Geographic Range, and Its Potential Status as a Soft Fruit Pest. *Journal of Integrated Pest Management*. 10: 20. DOI: [10.1093/jipm/pmz018](https://doi.org/10.1093/jipm/pmz018)
- R Core Team. 2024. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. <<https://www.R-project.org/>>
- Rakes LM, Delamont M, Cole C, Yates JA, Blevins LJ, Hassan FN, Bergland AO, Erickson PA. 2023. A small survey of introduced *Zaprionus indianus* (Diptera: Drosophilidae) in orchards of the eastern United States. *Journal of Insect Science*. 23: 21. DOI: [10.1093/jisesa/iead092](https://doi.org/10.1093/jisesa/iead092)
- Ramniwas S, Kajla B, Parkash R. 2012. Extreme physiological tolerance leads the wide distribution of *Zaprionus indianus* (Diptera: Drosophilidae) in temperate world. *Acta Entomologica Sinica*. 55: 1295.
- Renkema JM, Miller M, Fraser H, Legare JP, Hallett RH. 2013. First records of *Zaprionus indianus* Gupta (Diptera: Drosophilidae) from commercial fruit field in Ontario and Quebec, Canada. *The Journal of the Entomological Society of Ontario*. 144: 125.
- Renkema JM, Iglesias LE, Bonneau P, Liburd OE. 2018. Trapping system comparisons for and factors affecting populations of *Drosophila suzukii* and *Zaprionus indianus* in winter-grown strawberry. *Pest Management Science*. 74: 2076. DOI: [10.1002/ps.4904](https://doi.org/10.1002/ps.4904)
- Roque F, Matavelli C, Lopes PHS, Machida WS, Von Zuben CJ, Tidon R. 2017. Brazilian Fig Plantations Are Dominated by Widely Distributed Drosophilid Species (Diptera: Drosophilidae). *Annals of the Entomological Society of America*. 110: 521. DOI: [10.1093/aesa/sax044](https://doi.org/10.1093/aesa/sax044)
- Silva NM, Fantinel CC, Valente VLS, Valiati VH. 2005. Population dynamics of the invasive species *Zaprionus indianus* (Gupta) (Diptera: Drosophilidae) in communities of drosophilids of Porto Alegre City, Southern of Brazil. *Neotropical Entomology*. 34: 363. DOI: [10.1590/S1519-566X2005000300002](https://doi.org/10.1590/S1519-566X2005000300002)
- Stein CP, Teixeira P, Novo JPS. 2003. Aspectos biológicos da mosca do figo, *Zaprionus indianus* Gupta, 1970 (Diptera: Drosophilidae). *Entomotropica*. 18: 219.
- Tidon R, Leite DF, Leao BFD. 2003. Impact of the colonisation of *Zaprionus* (Diptera, Drosophilidae) in different ecosystems of the Neotropical Region: 2 years after the invasion. *Biological Conservation*. 112: 299. DOI: [10.1016/S0006-3207\(02\)00322-1](https://doi.org/10.1016/S0006-3207(02)00322-1)
- Van Der Linde K, Steck GJ, Hibbard K, Birdsley JS, Alonso LM, Houle D. 2006. FIRST RECORDS OF ZAPRIONUS INDIANUS (DIPTERA: DROSOPHILIDAE), A PEST SPECIES ON COMMERCIAL FRUITS FROM PANAMA AND THE UNITED STATES OF AMERICA. *Florida Entomologist*. 89: 402. DOI: [10.1653/0015-4040\(2006\)89\[402:FROZID\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2006)89[402:FROZID]2.0.CO;2)
- Vilela CR. 1999. Is *Zaprionus indianus* Gupta, 1970 (Diptera, Drosophilidae) currently colonizing the Neotropical region?. *Drosophila Information Service*. 82: 37.
- Wickham H. 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag, New York. DOI: [10.32614/CRAN.package.ggplot2](https://doi.org/10.32614/CRAN.package.ggplot2)
- Wilke C. 2024. *cowplot: Streamlined Plot Theme and Plot Annotations for 'ggplot2'*. Rpackage version 1.1.3. DOI: [10.32614/CRAN.package.cowplot](https://doi.org/10.32614/CRAN.package.cowplot)

7/17/2025 - Open Access

Yassin A, Capy P, Madi Ravazzi L, Ogereau D, David JR. 2008. DNA barcode discovers two cryptic species and two geographical radiations in the invasive drosophilid *Zaprionus indianus*. *Molecular Ecology Resources*. 8: 491. DOI: [10.1111/j.1471-8286.2007.02020.x](https://doi.org/10.1111/j.1471-8286.2007.02020.x)

Funding: Startup funds from the University of Richmond

Supported by National Institutes of Health (United States) 1R15GM146208 to Priscilla A. Erickson.

Supported by U.S. National Science Foundation (United States) 2145688 to Alan O. Bergland.

Author Contributions: Weston J. Gray: conceptualization, data curation, formal analysis, investigation, methodology, visualization, writing - original draft, writing - review editing. Megan Stephenson: investigation, data curation, writing - review editing. Alan O. Bergland: funding acquisition, supervision, writing - review editing. Priscilla A. Erickson: conceptualization, investigation, methodology, writing - original draft, writing - review editing.

Reviewed By: Anonymous

History: Received June 18, 2025 **Revision Received** July 1, 2025 **Accepted** July 15, 2025 **Published Online** July 17, 2025 **Indexed** July 31, 2025

Copyright: © 2025 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Gray WJ, Stephenson M, Bergland AO, Erickson PA. 2025. Field Sampling of *Zaprionus indianus* in the Eastern United States in 2023-2024. *microPublication Biology*. [10.17912/micropub.biology.001707](https://doi.org/10.17912/micropub.biology.001707)